

Risk category	Designated Class 2 medicines
Medicinal effect classification	Cold Remedies
Product name	ストナジェルサイナスEX STONA GEL SINUS EX
Product features	<ul style="list-style-type: none"> ● It is a cold medicine that contains the belladonna total alkaloid, diphenylpyraline hydrochloride, which has an excellent effect on the runny nose of the cold. ● Acetaminophen is effective in lowering fever, relieving headache and other pain. ● Made with an easy-to-swallow soft capsule that contains liquid.
Precautions	<p>■ When not to use the product (If you do not follow these instructions, the current symptoms may worsen or adverse reactions/incidents are more likely to occur.)</p> <ol style="list-style-type: none"> 1. This product should not be taken by the following persons. <ol style="list-style-type: none"> (1) Those who have experienced allergic symptoms to this medication or any of its ingredients. (2) Patients who have experienced asthma from taking this drug, other cold remedies, antipyretic analgesics. (3) Children under 15 years old. 2. This drug should not be taken together with the following drugs: Other cold medicines, antipyretics/analgesics, sedatives, expectorants, internal medicine, etc. containing antihistamines (internal medicine for rhinitis, medicine for motion sickness, medicine for allergies, etc.), gastrointestinal analgesics and antispasmodics 3. After taking this drug, do not drive a car or operate machinery (symptoms such as sleepiness, blurred vision and abnormal brightness may occur). 4. Nursing women should either not take this medicine or stop nursing while taking this medicine. 5. Do not drink alcohol before/after taking this medicine 6. Do not overdose or use for extended periods <p>■ Consultation</p> <ol style="list-style-type: none"> 1. The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before administration. <ol style="list-style-type: none"> (1) Patients undergoing medical treatment from a physician or dentist. (2) Pregnant women or women suspected of being pregnant. (3) The elderly. (4) Patients who have experienced allergic symptoms associated with drugs, etc. (5) Persons with the following symptoms: hyperthermia, urination difficulty (6) Those who have been diagnosed with the following: Thyroid disease, diabetes, heart disease, hypertension, liver disease, kidney disease, gastric/duodenal ulcer, glaucoma, respiratory dysfunction, obstructive sleep apnea syndrome, obesity 2. If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation. Related parts: Symptoms <ul style="list-style-type: none"> ○ Skin: rash/redness, itching ○ Gastrointestinal system: nausea and vomiting, loss of appetite, feeling of discomfort in the stomach, stomachache, abdominal pain, fullness of stomach and abdomen, heartburn, diarrhea ○ Neuropsychiatric system: dizziness, headache, numbness ○ Urinary system: urination difficulty ○ Other: excessive temperature decrease, facial hot flashes, abnormal brightness, oedema <p>The following serious symptoms may occur in rare cases. In such cases, immediately seek medical aid. Symptom name: Symptoms</p> <ul style="list-style-type: none"> ○ Shock (anaphylaxis): Symptoms, such as itching of skin, urticaria, hoarseness, sneezing, itchy throat, breathing difficulties, palpitations, and clouding of consciousness may occur immediately after take. ○ Oculomucocutaneous syndrome (stevens-johnson syndrome), toxic epidermal necrolysis, acute generalised exanthematous pustulosis: hyperthermia, ocular hyperaemia, eye discharge, lip erosion, pain throat, widespread skin rash/redness, small pimples (small pustules) on reddened skin, general malaise, anorexia, etc. may persist or suddenly worsen. ○ Drug-induced hypersensitivity syndrome: Symptoms such as widespread redness of the skin, generalised rash, fever, malaise, and swelling of lymph nodes (neck, armpit, groin, etc.) may occur. ○ Hepatic function failure: symptoms, such as fever, itching, rash, jaundice (yellowing of skin and white of eyes) , brown urine, general malaise, loss of appetite, etc. may occur. ○ Kidney disorders: Symptoms, such as fever, rash, reduced urinary volume, general oedema, general malaise, arthralgia (painful joints) and diarrhea, etc. may occur. ○ Interstitial pneumonia: shortness and/or difficulties of breath when go upstairs or overwork, sudden dry cough and/or fever and its continuance. ○ Asthma: Symptoms, such as wheezing or hissing when breathing, and difficult breathing, etc. may occur. ○ Respiratory depression: May cause shortness of breath, breathing difficulties, etc. <ol style="list-style-type: none"> 3. The following symptoms may be observed after taking this drug. If these symptoms persist or worsen, discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation. constipation, dry mouth, drowsiness, blurred vision 4. When symptoms do not improve even after taking the medicine 5 to 6 times, stop taking this medicine and consult your physician, pharmacist or registered salesperson, being sure to take this document with you.
Indication	Relief of various symptoms of a common cold: running nose, stuffy nose, sneezing, sore throat, cough, sputum, chills , fever, headache, joint pain, and muscle pain.
Dosage and administration	Take the following amount of medicine within 30 minutes after a meal. [Age: Single dose: Number of times per day] Adults (over 15) : 2 capsules: 3x Under 15 years old: Do not take
Precautions of Dosage and Administration	[1] Strictly follow the prescribed dosage and administration. [2] How to Dispense Capsules Using a finger, push the convex portion of the PTP sheet containing capsules to break the aluminum foil on the back.(Can lead to unforeseen consequences such as piercing the esophageal mucosa if mistakenly swallowed.)
Ingredient and amount	In 6 capsules Belladonna Total Alkaloids: 0.6mg: Works on the parasympathetic nerve system to suppress runny nose. Diphenylpyraline hydrochloride: 4mg: Relieves symptoms such as runny nose, stuffy nose, and sneezing. Acetaminophen: 900mg: Effective against fever, chills, headache, sore throat, and joint pain. dl-Methylephedrine hydrochloride: 60mg: Relieves cough symptoms. Dihydrocodeine phosphate: 24mg: Relieves cough symptoms. Noscapine: 48mg: Relieves cough symptoms. Ambroxol hydrochloride: 45mg: Make it easy to eject phlegm Anhydrous caffeine: 75mg: Improves efficacy of antipyretic and analgesic components. Riboflavin: 12mg: Supplements vitamins needed during a cold.
Precautions relating to ingredient and amount	<ul style="list-style-type: none"> ● Urine may turn yellow when taking this product as it contains Vitamin B2 Phosphate Ester.
Additive	Lauryl sulfate Na, polysorbate 80, propylene glycol fatty acid ester, glycerin fatty acid ester, gelatin, succinated gelatin, glycerin, D-sorbitol, titanium oxide, paraben

Precautions for storage and handling	<p>[1] Store in a cool, low humidity place away from direct sunlight.</p> <p>[2] Keep out of reach of children.</p> <p>[3] Do not transfer the medicine to a different container. (It may lead to misuse or alter the quality of the drug.)</p> <p>[4] Do not take the product after its expiration date.</p>
Customer Service Center	<p>For inquiries about this product, please contact the store where you purchased it or the following: Customer Service Center [Sato Pharmaceutical Co., Ltd.] Tel: 03-5412-7393 Reception hours: 9:00-17:00 (excluding Saturdays, Sundays, and holidays) Telephone inquiries are only available in Japanese as a language.</p>
Manufacturing and sales company	<p>Sato Pharmaceutical Co., Ltd. 1-5-27 Motoakasaka, Minato-ku, Tokyo</p>
Dosage form	<p>Capsule</p>
Pharmaceutical classification	<p>Over the counter drug</p>
Version No.	<p>5</p>
Contact details for the Adverse Reaction Relief System	<p>Pharmaceuticals and Medical Devices Agency https://www.pmda.go.jp/kenkouhigai_camp/index.html Tel: 0120-149-931 (toll-free)</p>
Disclaimer on Multilingual Product Information	<p>◆ Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. ◆ It does not warrant that its contents and the product itself conform to laws and regulations in countries other than Japan.</p>