

Risk category	Designated Class 2 medicines
Medicinal effect classification	Formulation containing antihistamine as the principal agent
Product name	スラジンA SURAZIN A
Product features	It is an anti-allergic medicine that combines chlorpheniramine maleate with antihistaminic activity and dl-methyl ephedrine hydrochloride with vasoconstrictive action, and extract of inchinko-to which is effective for hives.
Precautions	■ When not to use the product (If you do not follow these instructions, the current symptoms may worsen or adverse reactions/incidents are more likely to occur.) - This drug should not be taken together with the following drugs: Other allergy medicines, internal medicines containing antihistamines (cold remedy, antitussive expectorant, oral medicine for rhinitis, motion sickness medicine, etc.) - After taking this drug, do not drive a car or operate machinery (sleepiness may occur) . - Do not overdose or use for extended periods ■ Consultation - The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before administration. (1) Patients undergoing medical treatment from a physician. (2) Pregnant women or women suspected of being pregnant. (3) Nursing women. (4) The elderly. (5) Patients who have experienced allergic symptoms associated with drugs, etc. (6) Persons with the following symptoms: Urination difficulty (7) Those who have been diagnosed with the following: Heart disease, hypertension, diabetes, glaucoma, thyroid disease - If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation. Related parts: Symptoms ○ Skin: rash/redness, itching ○ Gastrointestinal system: nausea/vomiting, loss of appetite ○ Urinary system: urination difficulty The following serious symptoms may occur in rare cases. In such cases, immediately seek medical aid. Symptom name: Symptoms ○ Aplastic anemia: Symptoms, such as bruises, nose bleeds, bleeding gums, fever, pale appearance of skin and mucosa, fatigue, palpitations, shortness of breath, becoming dizzy from feeling sick and blood in the urine, etc. may occur. ○ Agranulocytosis: Symptoms, such as acute high fever, chills and sore throat, etc. may occur. ○ Mesenteric phleboscrosis: Abdominal pain, diarrhea, constipation, abdominal bloating, etc. may occur repeatedly after long-term use. - The following symptoms may be observed after taking this drug. If these symptoms persist or worsen, discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation. Dry mouth, drowsiness - When symptoms do not improve even after taking the medicine 5 to 6 times, stop taking this medicine and consult your physician, pharmacist or registered salesperson, being sure to take this document with you.
Indication	Hives, eczema, itching, rash
Dosage and administration	Take the following single dose after meals. [Age: Single dose: Number of times per day] Adult (15 years old and over): 1tablets: 3 times Under 15 years old: Do not take
Precautions of Dosage and Administration	[1] Strictly follow the prescribed dosage and administration. [2] How to Dispense Tablets Using your finger, push the convex portion of the PTP sheet containing capsules to break the aluminum foil on the back.(Can lead to unforeseen consequences such as piercing the esophageal mucosa if mistakenly swallowed.)
Ingredient and amount	In 3 tablets Chlorpheniramine maleate: 12mg: Antihistamine action is effective for allergic diseases such as hives and eczema. dl-Methyl ephedrine hydrochloride: 36mg: It works on sympathetic nerves and contracts capillaries, thereby relieving itching caused by hives, eczema, etc. Inchinko-to extract: 160mg: It is an ingredient consisting of 3 kinds of herbal medicines (Inchinko, gardenia fruit, rhubarb) that have an effect on hives
Additive	Lactose, hydroxypropyl-cellulose, magnesium stearate, hypromellose, polyoxyethylene polyoxypropylene glycol, polyvinyl alcohol (partially saponified product), calcium hydrogen phosphate, talc, polydimethylsiloxane, silicon dioxide, titanium oxide, sucrose, povidone, red No. 3, yellow No. 5, carnauba
Precautions for storage and handling	[1] Store in a cool, low humidity place away from direct sunlight. [2] Keep out of reach of children. [3] Do not transfer the medicine to a different container. (It may lead to misuse or alter the quality of the drug.) [4] Do not take the product after its expiration date.
Customer Service Center	For inquiries about this product, please contact the store where you purchased it or the following: Customer Service Center [Sato Pharmaceutical Co., Ltd.] Tel: 03-5412-7393 Reception hours: 9:00-17:00 (excluding Saturdays, Sundays, and holidays) Telephone inquiries are only available in Japanese as a language.
Other	[Advice to get better quickly] [1] When hives occur with food, try not to take that food. Try to have a balanced diet, avoid hot spices, and eat easily digestible food. [2] Do not cover the affected area. Doing so may cause suppuration, etc. and worsen symptoms. [3] Take a lukewarm bath. Itching may become stronger when taking a hot bath. [4] Arrange your bowel movements. It is important to excrete the causative food. Also, be careful not to irritate the digestive system as gastrointestinal disorders such as diarrhea may cause hives.
Manufacturing and sales company	Sato Pharmaceutical Co., Ltd. 1-5-27 Motoakasaka, Minato-ku, Tokyo
Dosage form	Tablet
Pharmaceutical classification	Over the counter drug
Version No.	9
Contact details for the Adverse Reaction Relief System	Pharmaceuticals and Medical Devices Agency https://www.pmda.go.jp/kenkouhigai_camp/index.html Tel: 0120-149-931 (toll-free)
Disclaimer on Multilingual Product Information	◆ Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. ◆ It does not warrant that its contents and the product itself conform to laws and regulations in countries other than Japan.