

Pharmaceutical classification	Over the counter drug
Medicinal effect classification	Gastrointestinal Medicines
product name	イノセアプラス錠 INOSEA PLUS TABLETS
Product features	<ul style="list-style-type: none"> <li>● Formulated with sucralfate hydrate for protection and repair of the mucous membrane of an upset stomach, magnesium aluminometasilicate for neutralization of excess stomach acid, and scopolia extract for inhibition of stomach acid secretion, for efficacy against stomachache.</li> <li>● Formulated with choleric ursodeoxycholic acid and digestive enzymes that aid digestion, for high efficacy against overdrinking and overeating.</li> <li>● Formulated with the antifoaming agent dimethylpolysiloxane for efficacy against feeling of fullness in the abdomen and stomach.</li> </ul>
Indication	Stomachache, feeling of fullness in the stomach, feeling of discomfort in the stomach, heaviness in the stomach, heaviness in the chest, constricted feeling in the stomach (stomach heaviness), heartburn, belching (burping), nausea (retching, stomach retching, retching due to hangovers and overdrinking, sick feeling, feeling of sickness), vomiting, loss of appetite (anorexia), excessive drinking (overdrinking), excessive eating (overeating), promoting digestion, indigestion, heaviness in the chest, and feeling of fullness in the stomach and abdomen due to indigestion, and hyperacidity
Dosage and administration	Take the following dose once a day after or between meals.  [Age: single dose: number of daily doses] Adult (over 15 years old): 4 tablets: 3x Under 15 years: Do not take.
Precautions of Dosage and Administration	Strictly follow the prescribed dosage and administration.
Ingredient and amount	<p>In 12 tablets</p> <p>Sucralfate hydrate : 1500mg : Selectively binds to lesions in the gastric mucosa for protection and repair of stomach lining, making it effective for stomach pain.</p> <p>Magnesium aluminometasilicate : 900mg : Neutralizes excess stomach acid and works to protect stomach lining.</p> <p>Dimethylpolysiloxane : 78.95mg : Works as an anti-foaming agent to reduce the volume of stomach content, making it effective against bloated stomach.</p> <p>Scopolia extract : 30mg : Acts on the nerves that regulate stomach movement, inhibits excessive stomach acid secretion and alleviates stomachache.</p> <p>Dried atracylodes lancea extract : 60mg : Acts on the nerves that regulate stomach movement, inhibits excessive stomach acid secretion and alleviates stomachache.</p> <p>Diasmen SS : 60mg : A powerful starch digestive enzyme to aid digestion.</p> <p>Lipase AP6 : 60mg : A fat digestive enzyme to aid digestion.</p> <p>Ursodeoxycholic acid : 30mg : A choleric discovered in the gallbladder of a bear shows excellent efficacy against upset stomach, nausea, and drunkenness.</p>
Precautions	<p>■When not to use the product (If you do not follow these instructions, the current symptoms may worsen or adverse reactions are more likely to occur.)</p> <ol style="list-style-type: none"> <li>1. This product should not be taken in the following persons. Patients undergoing dialysis.</li> <li>2. This medicine should not be taken together with the following medicines: Pain relief/antispasmodic medicine for the stomach</li> <li>3. For those who are lactating, do not take this medicine or avoid lactating while using this medicine.(The medicine may be transferred to the breast milk and accelerate the pulse of the infant. )</li> <li>4. Do not take this medicine for a long time.</li> </ol> <p>■Consultation</p> <ol style="list-style-type: none"> <li>1. The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before administration. (1)Patients undergoing medical treatment from a physician (2)Pregnant women or women suspected of being pregnant (3)The elderly (4)Patients who have experienced allergic symptoms associated with drugs, etc. (5)Persons with the following symptoms urination difficulty (6)Persons diagnosed as having the following kidney disease, heart disease, glaucoma</li> </ol>

	<p>2. If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this product to your physician, pharmacist, or registered salesperson for a consultation.</p> <p>&lt;Related parts : Symptoms&gt; Skin : rash/redness, itching</p> <p>3. The following symptoms may be observed after taking this drug. If these symptoms persist or worsen, discontinue the use of this drug, and show this product to your physician, pharmacist, or registered salesperson for a consultation.</p> <p>dry mouth, constipation, diarrhea</p> <p>4. When symptoms do not improve even after taking this medicine for about 2 weeks, stop taking this medicine and consult a physician, pharmacist or registered salesperson, being sure to take this product with you.</p> <p>&lt;Other notes&gt; Some may have trouble lactating.</p>
Additive	Cellulose, CMC-Ca, lactose, macrogol, povidone, magnesium stearate, silicic acid anhydride, calcium carbonate, cross CMC-Na, fragrances (including l-menthol)
Precautions for storage and handling	<p>(1)Store in a cool, low humidity place away from direct sunlight.</p> <p>(2)Keep out of reach of children.</p> <p>(3) Do not transfer the medicine to a different container. (It may lead to misuse or alter the quality of the drug. )</p> <p>(4) Do not take the product after its expiration date.</p>
Other	<p>&lt; Advice to get better quickly &gt;</p> <ul style="list-style-type: none"> <li>●Refrain from consuming liquor, cigarettes, sour foods, spices (pepper, chili pepper, etc.),coffee, and tea to the extent possible.</li> <li>●To avoid stressing the stomach, consume foods that are easy to digest.</li> <li>●Also stay conscious of mental aspects, such as irritation and nervousness.</li> </ul>
Manufacturing and sales company	Sato Pharmaceutical Co., Ltd. 1-5-27 Moto Akasaka, Minato-ku, Tokyo
Risk category	Category 2 OTC medicine
Disclaimer on Multilingual Product Information	<ul style="list-style-type: none"> <li>• This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.</li> <li>• Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.</li> <li>• Multilingual product information is a tentative translation by Our Company, and may be modified or altered without notice.</li> <li>• Our Company assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.</li> </ul>