

Pharmaceutical classification	Over the counter drug
Medicinal effect classification	Agent for arteriosclerosis
product name	ラングロン LUNGRON
Product features	<ul style="list-style-type: none"> ● Lowers the cholesterol level in the blood and is effective for improving high serum cholesterol. Cholesterol in the blood is an important element that is necessary for making cells and hormones in the body, but if there is too much in the blood, it accumulates in the walls of blood vessels and causes arteriosclerosis. ● Riboflavin butyrate prevents cholesterol from being produced in the body, and has the effect of draining the produced cholesterol, which is effective in improving high serum cholesterol, out of the body. ● A red-purple soft capsule which contains riboflavin butyrate dissolved in vegetable oil
Indication	Improvement of hypercholesterolemia,
Dosage and administration	<p>Take the following dose twice a day.</p> <p><Age: single dose: number of daily doses></p> <p>Adult (15 years old and over): 1 capsules: 2 times</p> <p>Under 15 years: Do not take.</p>
Precautions of Dosage and Administration	<p>(1)Strictly follow the prescribed dosage and administration.</p> <p>(2)Dietary therapy is important for the improvement of high serum cholesterol, so please continue dietary therapy even if you take this drug.</p>
Ingredient and amount	<p>In 2 capsules</p> <p>Riboflavin butyrate : 60 mg : It suppresses cholesterol synthesis in the liver and improves blood lipid metabolism by excretion or catabolism of blood cholesterol. It also increases HDL (good) cholesterol. These actions are effective in improving high cholesterol levels in the blood.</p>
Precautions	<p>■Consultation</p> <p>1. The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before administration.</p> <p>(1)Patients undergoing medical treatment from a physician</p> <p>(2)Patients who have experienced allergic symptoms associated with drugs, etc.</p> <p>2. If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation.</p> <p><Related parts : Symptoms></p> <p>Skin: rash/redness, itching</p> <p>Gastrointestinal system: nausea, feeling of discomfort in the stomach, heartburn</p> <p>3. The following symptoms may be observed after taking this drug. If these symptoms persist or worsen , discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation.</p> <p>diarrhea</p> <p>4. When symptoms do not improve even after taking this medicine for a while, stop taking this medicine and consult a physician, pharmacist or registered salesperson, being sure to take this instruction leaflet with you.</p>
Additive	Medium chain fatty acid triglyceride, d-δ-tocopherol, gelatin, glycerin, D-sorbitol, paraben, titanium oxide, red No. 106, yellow No. 5
Precautions for storage and handling	<p>(1) Store in a cool, low humidity place away from direct sunlight.</p> <p>(2)Keep out of reach of children.</p> <p>(3)Do not transfer the medicine to a different container. (It may lead to misuse or alter the quality of the drug.)</p> <p>(4)Do not take the product after its expiration date.</p> <p>(5)Capsules easily absorb moisture, so be careful not to touch them with wet hands.</p>
Other	<p>< People with high cholesterol levels ...></p> <p>It is important to take care of your daily meals.</p> <p>1. Try to eat a good balance of animal fat (contained in meat and butter) and vegetable fat (contained in olives and soy).</p> <p>2. Try to eat brown rice, soy, etc. that contain a lot of vitamin E.</p> <p>3. Try to moderate eggs, eels, squid, liver, etc. high in cholesterol.</p> <p>4. Try to moderate salt and sugars.</p> <p>5. Try to eat a lot of dietary fiber.</p> <p>6. If you take more calories than you need before going to bed, it will make extra cholesterol, so try to refrain from eating at night.</p>
Manufacturing and sales company	<p>Sato Pharmaceutical Co., Ltd.</p> <p>1-5-27 Moto Akasaka, Minato-ku, Tokyo</p>

Risk category	Category 3 OTC medicine
Disclaimer on Multilingual Product Information	<ul style="list-style-type: none">• This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.• Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.• Multilingual product information is a tentative translation by Our Company, and may be modified or altered without notice.• Our Company assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.